

Name: _____

Date: _____

UNIT 7


LESSON 7

AIM: HOW DO WE DETERMINE THE LENGTH BETWEEN TWO POINTS?


Finding length given two coordinates (x, y) .

_____ FORMULA $d =$


1. The coordinates of point A are $(0,3)$, the coordinates of point B are $(4,0)$ and the coordinates of point C are $(0,0)$. What is the length of \overline{AB} ?




2. The coordinates of point A are $(4,8)$, the coordinates of point B are $(4,2)$ and the coordinates of point C is $(1,2)$. What is the length of \overline{AC} , in simplest radical form?



3. The point $(5,4)$ lies on a circle. What is the length of the **diameter** of this circle if the center is located at $(3,2)$ in simplest radical form?



4. The coordinates of rectangle ABCD are $A(0,2)$, $B(4,8)$, $C(7,6)$ and $D(3,0)$. Show that the diagonals are equal in length.



5. Given triangle DEF with coordinates $D(2, 3)$, $E(7, 5)$ and $F(4, -2)$.
- Find the length of each side of triangle DEF

b. Based on part a, what type of triangle is DEF? Explain.

6. The coordinates of the vertices of $\triangle DEF$ are $D(-2,0)$, $E(4,0)$, and $F(1,3\sqrt{3})$
- Find the length of DE, EF, and FD

b. Is $\triangle EDF$ equilateral? Justify your answer.

REVIEW:

1. Line segment RW has endpoints $R(-4,5)$ and $W(6,20)$. Point P is on RW such that $RP:PW$ is 2:3. What are the coordinates of point P ?

1) (2,9)

2) (0,11)

3) (2,14)

4) (10,2)

<p>2. A parallelogram must be a rectangle if its diagonals</p> <ol style="list-style-type: none">1) bisect each other.2) bisect the angles to which they are drawn.3) are perpendicular to each other.4) are congruent.	<p>3. Which statements describe the properties of a trapezoid?</p> <ol style="list-style-type: none">1) The bases are parallel.2) The diagonals are congruent.3) The opposite angles are congruent.4) The base angles are congruent.
<p>4. Which of the following reasons is valid for proving a quadrilateral is a parallelogram?</p> <ol style="list-style-type: none">1) Diagonals bisect angles2) All sides are congruent3) One pair of opposite sides are parallel4) One pair of opposite sides are both parallel and congruent	<p>5. Which of the following reasons is NOT valid for proving a parallelogram is a rhombus?</p> <ol style="list-style-type: none">(1) Diagonals bisect angles(2) All sides are congruent(3) Diagonals are congruent(4) Diagonals are perpendicular
<p>6. Which of the following reasons is valid for proving a parallelogram is a rectangle?</p> <ol style="list-style-type: none">(1) Diagonals bisect angles(2) Both pairs of opposite sides are congruent(3) Diagonals are congruent(4) Diagonals are perpendicular	<p>7. The diagonals of a quadrilateral are congruent but do not bisect each other. This quadrilateral is</p> <ol style="list-style-type: none">1) an isosceles trapezoid2) a parallelogram3) a rectangle4) a rhombus

Name: _____

Date: _____

UNIT 7

LESSON 7 HOMEWORK

1. The center of circle O has coordinates $(6, 4)$. If circle O passes through $(-9, -4)$, what is the length of its diameter?

2. The endpoints of one side of a regular octagon are $(-1, 4)$ and $(2, 3)$. What is the perimeter of the octagon?


HINT: Regular polygons have equal sides and equal angles!

- 1) $\sqrt{10}$
- 2) $8\sqrt{10}$
- 3) $8\sqrt{2}$
- 4) $64\sqrt{2}$

3. The vertices of square $RSTV$ have coordinates $R(-1, 5)$, $S(-3, 1)$, $T(-7, 3)$, and $V(-5, 7)$. What is the perimeter of $RSTV$?

- 1) $\sqrt{20}$
- 2) $\sqrt{40}$
- 3) $4\sqrt{20}$
- 4) $4\sqrt{40}$

4. If $\triangle MNP$ has vertices at $M(-5, -7)$, $N(7, -2)$ and $P(2, 10)$. Is $\triangle MNP$ isosceles? Explain your answer.



5. Directed segment \overrightarrow{AB} is drawn from $A(-7, -4)$ to $B(0, 10)$. Find point C that partition \overrightarrow{AB} in the ratio 5:2.